

Similar radiation mechanism in gamma-ray bursts and blazars: evidence from two luminosity correlations

F. Y. Wang^{1,2}, S. X. Yi^{1,2} and Z. G. Dai^{1,2}

¹*School of Astronomy and Space Science, Nanjing University, Nanjing 210093, China*

²*Key Laboratory of Modern Astronomy and Astrophysics (Nanjing University), Ministry of Education, Nanjing 210093, China*

ABSTRACT

Active galactic nuclei (AGNs) and gamma-ray bursts (GRBs) are powerful astrophysical events with relativistic jets. In this Letter the broadband spectral properties are compared between GRBs and the well-observed blazars. The distribution of GRBs are consistent with the well-known blazar sequence including the $\nu L_\nu(5\text{GHz}) - \alpha_{\text{RX}}$ and $\nu L_\nu(5\text{GHz}) - \nu_{\text{peak}}$ correlations, where α_{RX} is defined as the broadband spectral slope in radio-to-X-ray bands, and ν_{peak} is defined as the spectral peak frequency. Moreover, GRBs occupy the low radio luminosity end of these sequences. These two correlations suggest that GRBs could have a similar radiation process with blazars both in the prompt emission and afterglow phases, i.e., synchrotron radiation.

Subject headings: gamma-ray burst: general - BL Lacertae objects: general - methods: statistical

1. Introduction

Gamma-ray bursts (GRBs) and active galactic nuclei (AGNs) are both powered by relativistic jets from accreting black holes (Gehrels et al. 2009; Urry & Padovani 1995). The central engines of GRBs are argued to be stellar-mass black holes (Woosley 1993) and for AGNs the central engines are supermassive black holes. GRBs are most powerful explosions with isotropic-equivalent energy $E_{\text{iso}} \sim 10^{50} - 10^{55}$ erg in the universe (Zhang 2011), and can be detected out to very high-redshift universe (Lamb & Reichart 2000; Wang et al. 2012). So GRBs can probe high-redshift universe, including dark energy (Dai, Liang & Xu 2004;

E-mail: fayinwang@nju.edu.cn (FYW); dzg@nju.edu.cn (ZGD)

Schaefer 2007; Wang & Dai 2011). Blazars include two subtype of AGNs, i.e., flat-spectrum radio quasars (FSRQs) and BL Lac objects (BL Lacs). A subclass of AGNs, e.g., super-Eddington accreting supermassive black holes, are also proposed to be a standard candle (Wang et al. 2013).

The radiation mechanism of blazars is well constrained. The spectral energy distribution (SED) of blazars is well understood, including the low-energy (infrared-soft X-ray) bump and the high-energy (MeV-GeV) bump. The synchrotron radiation can account for the low-energy peak, while the MeV-GeV peak is produced by inverse Compton radiation. But for GRBs, the radiation mechanism for the prompt emission is still highly debated. The spectrum of prompt emission can be modeled by the “Band function” (Band et al. 1993), whose origin is still unknown (but see Lucas Uhm & Zhang 2013). Some studies (Mészáros et al. 1994; Daigne & Mochkovitch 1998) proposed that synchrotron radiation is the leading mechanism. Other mechanisms are also proposed (Pe’er et al. 2006; Rees & Mészáros 2005; Beloborodov 2010). The radiation mechanism of afterglows is well understood (Sari et al. 1998). The observed afterglow radiation is well explained by synchrotron radiation (Sari et al. 1998; Panaitescu & Kumar 2001).

Some studies (Zhang 2007; Wang & Dai 2013; Wang et al. 2014) have proposed that the mechanisms in different scale outflow or jet systems may be the same. Some works have been done on comparison between GRBs and AGNs. Wang & Wei (2011) compared the spectral properties of blazars and optically bright GRB afterglows, and found that GRB afterglows have the same radiation mechanism as BL Lac objects. A similar correlation of the synchrotron luminosity and Doppler factor between GRBs and AGNs has been found (Wu et al. 2011). Nemmen et al. (2012) suggested that the relativistic jets in AGNs and GRBs have a similar energy dissipation efficiency. Ma et al. (2014) extended the analysis of Nemmen et al. (2012) by adding X-ray binaries and low-luminosity AGNs. Wang & Dai (2013) found that the GRB X-ray flares and solar X-ray flares have similar distributions, which indicate that the X-ray flares of GRBs are due to a magnetic reconnection process. These similar distributions also exist in X-ray flares from black hole systems with $10^6 - 10^9 M_\odot$ (Wang et al. 2014). Zhang et al. (2013) found that the prompt emission of GRBs may be produced by magnetic dominated jets.

In this Letter we compare the broadband spectral properties of GRBs and blazars, including the $\nu L_\nu(5\text{GHz}) - \alpha_{RX}$ and $\nu L_\nu(5\text{GHz}) - \nu_{\text{peak}}$ correlations, where α_{RX} is the radio-to-X-ray spectral slope. For a GRB, ν_{peak} is the peak frequency of νf_ν spectrum of prompt emission, while ν_{peak} is the low peak of νf_ν spectrum for a blazar. The aim of this Letter is to explore a possible similarity in radiation mechanism between GRBs and blazars. this Letter is organized as follows. In section 2, we present the sample of blazars and GRBs. The

fitting results are given in section 3. Section 4 gives conclusions and discussions.

2. Samples

Chandra & Frail (2012) compiled a sample of GRB radio afterglow observations from 1997 to 2011. This catalog consists of 304 GRBs with radio observations. We select 43 GRBs with redshift measurements. The sample is listed in Table 1. For each GRB, the name of GRB and redshift are presented in Columns 1 and 2, respectively. X-ray flux measured in 0.3-10 keV energy band is given in Column 3 observed by Swift at the time of Column 6. For bursts observed by BASTE, the energy range is between 2 keV and 10 keV. We use the typical GRB spectrum to convert the flux to 1 keV. The observed radio flux (Column 4) and frequency (Column 5) at the time of Column 6 are also provided. Column 7 is the jet opening angle. For GRBs without opening angle determination, we assume a typical value 5 degree. The derived collimation-corrected radio luminosity at 5 GHz is given in Column 8. Column 9 gives E_{peak} , which is the peak energy of the prompt spectrum. The parameters from Column 2 to Column 7 are taken from Chandra & Frail (2012). We use the value of E_{peak} from Wang, Qi & Dai (2011). In the calculation, we use the peak frequency in the GRB rest frame. The collimation-corrected radio luminosity is calculated as

$$\nu L_\nu(5\text{GHz}) = 4\pi d_L^2 \nu f_\nu F_{\text{beam}}(1+z)^{-\alpha-1}, \quad (1)$$

where d_L is the luminosity distance, f_ν is radio flux at 5 GHz, $F_{\text{beam}} = 1 - \cos \theta_j$ is the beaming factor and $\alpha = 1/3$ is the spectral slope (Sari et al. 1998). We adopt $\alpha = 1/3$ in the slow cooling case. The observed radio flux is converted to flux at 5 GHz using spectral slope $\alpha = 1/3$. In this work, we assume the cosmological parameters: $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_m = 0.3$, and $\Omega_\Lambda = 0.7$.

We use the spectral properties of blazars from Fossati et al. (1998). This sample consists of all the parameters that we require, including redshift, X-ray flux at 1 keV, radio flux at 5 GHz, and synchrotron peak frequency. Fossati et al. (1998) found a power spectral sequence for the blazars despite the difference in the continuum shapes among different sub-classes of blazars. This sequence indicates that the radio luminosity is anti-correlated with the synchrotron peak. A plausible interpretation is that relativistic jets radiate via synchrotron and inverse Compton processes if the physical parameters (i.e. magnetic field) vary with luminosity.

3. Results

3.1. $\nu L_\nu(5\text{GHz}) - \nu_{\text{peak}}$ correlation in blazars and GRBs

The radio luminosities at 5 GHz are anti-correlated with the synchrotron peaks of blazars, as found by Fossati et al. (1998). This correlation has not been studied in GRBs so far. We investigate this correlation in GRBs for the first time. Figure 1 shows the $\nu L_\nu(5\text{GHz}) - \nu_{\text{peak}}$ correlation of blazars and GRBs. The black and open dots represent blazars and GRBs, respectively. There is a tight correlation between $\nu L_\nu(5\text{GHz})$ and ν_{peak} as expected from the blazar sequence (Fossati et al. 1998). The correlation coefficient is $r = 0.83$ at a significance level $p < 10^{-4}$ from Spearman rank-order statistical test. From this figure, we can see that the GRBs occupy the low-luminosity region of this correlation. For both blazars and GRBs, the best fitting result is

$$\log \nu L_\nu(5 \text{ GHz}) = (-0.91 \pm 0.03) \log \nu_{\text{peak}} + 56.65 \pm 0.46. \quad (2)$$

The correlation coefficient is improved to $r = 0.94$ with probability $p < 10^{-4}$. The correlation coefficient has an obvious enhancement after adding the GRB sample. GRB 060218 may deviate from this correlation. The possible reason is that this GRB usually called X-ray flash has a low peak energy (Soderberg et al. 2006; Pian et al. 2006).

3.2. $\nu L_\nu(5\text{GHz}) - \alpha_{\text{RX}}$ correlation in blazars and GRBs

The broad-band spectral slope α_{RX} is also correlated with luminosity at 5 GHz in blazars (Fossati et al. 1998). We also investigate this correlation in GRBs. The broad-band spectral slope α_{RX} is defined as

$$\alpha_{\text{RX}} = -\frac{\log(f_{\nu_R})/\log(f_{\nu_x})}{\log(\nu_R/\nu_x)}, \quad (3)$$

where $\nu_R = 5 \text{ GHz}$, and $\nu_x = 1 \text{ keV}$. The X-ray flux f_{ν_x} at 1 keV can be obtained as follows. From Column 3 of Table 1, X-ray flux measured in 0.3-10 keV energy range can be obtained. The X-ray flux usually evolves as $\nu^{-p/2}$ (Sari et al. 1998), with $p = 2.2$ is the power-law index of accelerated electrons distribution. The X-ray flux in 0.3-10 keV is $\int_{0.3\text{keV}}^{10\text{keV}} a\nu^{-p/2} d\nu = F_X$. After obtaining the value of a , the flux at 1 keV can be derived.

Figure 2 shows a correlation between the spectral slope α_{RX} and radio luminosity at 5 GHz for blazars (black dots) and GRBs (open dots). The correlation coefficient is $r = 0.82$ with probability $p < 10^{-4}$ using Spearman rank-order statistical test for blazars. After combining GRBs and blazars, the fitting result is

$$\alpha_{\text{RX}} = (0.084 \pm 0.006) \log(\nu L_\nu(5\text{GHz})) + (-2.87 \pm 0.24). \quad (4)$$

The Spearman’s rank correlation coefficient is $r = 0.85$ with probability $p < 10^{-4}$. GRBs also occupy the the low-luminosity end of this correlation.

4. Conclusions and Discussions

The physics of GRBs are poorly understood, i.e., the radiation mechanism of prompt emission, the value of Lorentz factor, jet composition, and central engine (Zhang 2011). Wang & Dai (2013) found similar frequency distributions between X-ray flares of GRBs and solar X-ray flares, which may indicate the magnetically dominated jets in GRBs. In this Letter we compile 43 GRBs with well X-ray and radio observations. Two new correlations between GRBs and blazars may provide a new clue as to the radiation mechanism of GRB prompt emission and afterglows. For example, our clear $\nu L_\nu(5\text{GHz}) - \alpha_{\text{RX}}$ and $\nu L_\nu(5\text{GHz}) - \nu_{\text{peak}}$ correlations suggest that the radiation mechanism of GRBs in prompt and afterglow phases and blazars is similar, namely, synchrotron radiation. Moreover, GRBs are occupy the low-luminosity region of these correlations.

In this Letter, we use the radio luminosities during the GRB afterglow phase. Although some models predict that bright radio emission may be generated within about 10 s of the initial explosion of a GRB (Usov & Katz 2000; Sagiv & Waxman 2002; Shibata et al. 2011), but a detection of prompt radio emission has some impediments, such as scattering (Macquart 2007; Lyubarsky 2008). Bannister et al. (2012) have searched for prompt radio emission from nine GRBs at 1.4 GHz, and found single dispersed radio pulses with significance $> 6\sigma$ in a few minutes following two GRBs. Unfortunately, the probability of GRB origin is only 2%. There has been no confirmed evidence for detection of GRB prompt radio emission up to now. So we use the radio emission of an afterglow in this Letter. Theoretically, internal shocks produce the GRB prompt emission, and external shocks produce the afterglow emission. The radiation mechanism of an afterglow is well constrained, i.e., synchrotron emission. But the prompt emission related to ν_{peak} is not well understood. If some similar correlations between $\nu L_\nu(5\text{GHz})$ and ν_{peak} exist in GRBs and blazars, the radiation mechanism of prompt emission is argued as synchrotron emission. Until now, the location of blazar gamma-ray emission regions are still uncertain (Marscher et al. 2010), since some theories locate blazar gamma-ray emission regions close to the black hole/accretion disk (Blandford & Levinson 1995) while the others place them at parsec scales in the radio jet (Jorstad et al. 2001). The X-ray emission from kiloparsec-scale blazar jets has been observed (Harris & Krawczynski 2006). Meanwhile, the radio emission region of blazars spans kiloparsec scale. Kharb et al. (2010) found that a few blazars exhibit only radio core emission. But the X-ray emission region and radio emission region do not fully overlap. So the high-energy

and radio emissions of blazars also originate from different regions. So a comparison of the correlation $\nu L_\nu(5\text{GHz}) - \nu_{\text{peak}}$ between GRBs and the blazar sequence is reasonable. Liang et al. (2004) found that the peak energy ν_{peak} evolves with isotropic-equivalent luminosity from the time-resolved spectra. They also found that the $L_{\text{iso}} - \nu_{\text{peak}}$ correlation also holds for time-resolved spectra and time-integrated spectra. Ghirlanda et al. (2010) studied the time-resolved spectra of Fermi GRBs. The peak energy ν_{peak} correlates with the luminosity within individual bursts (Ghirlanda et al. 2010). Moreover, the time-resolved $L_{\text{iso}} - \nu_{\text{peak}}$ correlation is very similar for all the bursts and has a slope similar to the correlation defined by the time-integrated spectra of different bursts detected by several different satellites. The time-integrated value of ν_{peak} is widely used in luminosity correlations of GRBs, for example $\nu_{\text{peak}} - E_{\text{iso}}$ (Amati et al. 2002), $\nu_{\text{peak}} - L_{\text{iso}}$ (Yonetoku et al. 2004) and $\nu_{\text{peak}} - E_{\gamma, \text{jet}}$ (Ghirlanda et al. 2004) correlations. So if some correlation is due to a similar physical mechanism, this correlation holds no matter whether the time-resolved or time-integrated values are used.

Fossati et al. (1998) found that ν_{peak} is anti-correlated with the synchrotron peak luminosity for blazars. For GRBs, Liang et al. (2004) found that ν_{peak} is positively correlated with the isotropic-equivalent luminosity (about total luminosity). But the luminosity in the correlation for blazars is the synchrotron peak luminosity, not the total luminosity. Because there are two peaks in a blazar spectral energy distribution and there is no correlation between E_{iso} and $\nu L_\nu(5\text{GHz})$ in GRBs (Chandra & Frail 2012), from our simple analysis above we cannot conclude that GRBs have a different $\nu L_\nu(5\text{GHz}) - \nu_{\text{peak}}$ correlation compared with that for blazars. In this paper, we find that GRBs occupy the low radio luminosity end of the blazar sequence, which is similar to that of Wang & Wei (2011).

We thank the referee for detailed and very constructive suggestions that have allowed us to improve our manuscript. We acknowledge helpful discussions with Y. C. Zou and X. F. Wu. This work is supported by the National Basic Research Program of China (973 Program, grant No. 2014CB845800) and the National Natural Science Foundation of China (grants 11373022, 11103007, and 11033002).

REFERENCES

- Amati, L., Frontera, F., Tavani, M., et al. 2002, *A&A*, 390, 81
- Band, D., Matteson, J., Ford, L., et al. 1993, *ApJ*, 413, 281
- Bannister, K. W., Murphy, T., Gaensler, B. M. & Reynolds, J. E., 2012, *ApJ*, 757, 38

- Blandford, R. D., & Levinson, A. 1995, *ApJ*, 441, 79
- Chandra, P. & Frail, D. A., 2012, *ApJ*, 746, 156
- Dai, Z. G., Liang, E. W., & Xu, D. 2004, *ApJ*, 612, L101
- Daigne, F., & Mochkovitch, R. 1998, *MNRAS*, 296, 275
- Beloborodov, A. M., 2010, *MNRAS*, 407, 1033
- Fardal, M. A., Katz, N., Weinberg, D. H. & Dave, R., 2007, *MNRAS*, 379, 985
- Ferguson, H. C., Dickinson, M., & Papovich, C., 2002, *ApJ*, 569, L65
- Fossati, G., Maraschi, L., Celotti, A., Comastri, A., & Ghisellini, G. 1998, *MNRAS*, 299, 433
- Friedmann, A. S., & Bloom, J. S. 2005, *ApJ*, 627, 1
- Gehrels, N., Ramirez-Ruiz, E. & Fox, D. B., 2009, *ARA&A*, 47, 567
- Ghirlanda, G., Ghisellini, G. & Lazzati, D., 2004, *ApJ*, 616, 331
- Ghirlanda, G., Nava, L. & Ghisellini, G., 2010, *A&A*, 511, A43
- Harris, D. E., & Krawczynski, H., 2006, *ARA&A*, 44, 463
- Jorstad, S. G., Marscher, A. P., Mattox, J. R., et al. 2001, *ApJ*, 556, 738
- Kharb, P., Lister, M. L. & Cooper, N. J., 2010, *ApJ*, 710, 764
- Lamb, D. Q., & Reichart, D. E. 2000, *ApJ*, 536, 1
- Liang, E. W., Dai, Z. G. & Wu, X. F., 2004, *ApJ*, 606, L29
- Lucas Uhm, Z., & Zhang, B., 2013, arXiv: 1303.2704
- Lyubarsky, Y. 2008, *ApJ*, 682, 1443
- Ma, R. Y., Xie, F. G., & Hou, S. J., 2014, *ApJL*, 780, L14
- Macquart, J. P., 2007, *ApJL*, 658, L1
- Marscher, A. P., Jorstad, S. G., Larionov, V. M., et al. 2010, *ApJ*, 710, L126
- Mészáros, P., Rees, M. J., & Papathanassiou, H. 1994, *ApJ*, 432, 181

- Nemmen, R. S., Georganopoulos, M., Guiriec, S., et al. 2012, *Sci*, 338, 1445
- Panaiteanu, A., & Kumar, P., 2001, *ApJ*, 560, L49
- Pe’er, A., Mészáros, P., & Rees, M. J., 2006, *ApJ*, 642, 995
- Pian, E., Mazzali, P. A., Masetti, N., et al., 2006, *Nature*, 442, 1011
- Rees, M. J., & Mészáros, P., 2005, *ApJ*, 628, 847
- Sagiv, A., & Waxman, E. 2002, *ApJ*, 574, 861
- Sari, R., Piran, T., & Narayan, R. 1998, *ApJ*, 497, L17
- Schaefer, B. E., 2007, *ApJ*, 660, 16
- Soderberg, A. M., Kulkarni, S. R., Nakar, E., et al., 2006, *Nature*, 442, 1014
- Urry, C. M., & Padovani, P. 1995, *PASP*, 107, 803
- Usov, V. V., & Katz, J. I. 2000, *A&A*, 364, 655
- Wang, F. Y., Bromm, V., Greif, T. H., Stacy, A., Dai, Z. G., Loeb, A. & Cheng, K. S. 2012, *ApJ*, 760, 27
- Wang, F. Y., & Dai, Z. G., 2013, *Nature Phys.*, 9, 465
- Wang, F. Y., Dai, Z. G., & Yi, S. X., 2014, submitted
- Wang, F. Y., & Dai, Z. G. 2011, *A&A*, 536, A96
- Wang, F. Y., Qi, S., & Dai, Z. G., 2011, *MNRAS*, 415, 3423
- Wang, J., & Wei, J. Y. 2011, *ApJL*, 726, L4
- Wang, J. M., Du, P., Valls-Gabaud, D., et al. 2013, *PRL*, 110, 081301
- Woosley, S. E., *ApJ*, 1993, 405, 273
- Wu, Q., Zou, Y. C., Cao, X., Wang, D. X., & Chen, L. 2011, *ApJL*, 740, L21
- Yonetoku, D., Murakami, T., Nakamura, T. et al., 2004, *ApJ*, 609, 935
- Zhang, B. 2011, *CRPhy*, 12, 206
- Zhang, J., Liang, E. W., Sun, X. N., et al., 2013, *ApJL*, 774, L5

Zhang, S. N. 2007, HiA, 14, 41

Table 1. GRB data

GRB	z	F_X	F_R	Frequency	T_{peak}	θ_j	L_R	E_p
		$10^{-13} \text{erg cm}^{-2} \text{s}^{-1}$	μJY	GHz	days		erg s^{-1}	keV
970508	0.835	1.2	780 $\pm$ 13	4.86	31.4	20.93	3.83E39	389 $\pm$ 40
970828	0.958	8.0	144 $\pm$ 31	8.46	4.0	5.15	6.70E37	298 $\pm$ 30
980329	3	...	171 $\pm$ 14	4.86	22.9	3.34	1.82E38	134 $\pm$ 20
980425	0.009	3.8	38362 $\pm$ 337	4.80	18.1	5.00	1.28E36	150 $\pm$ 35
980703	0.966	4.2	1055 $\pm$ 30	4.86	4.6	14.82	3.42E39	254 $\pm$ 25
990510	1.619	2.5	255 $\pm$ 34	8.46	1.6	3.36	1.27E38	126 $\pm$ 10
010222	1.477	1.2	93 $\pm$ 25	8.46	6.8	3.02	3.21E37	309 $\pm$ 12
011121	0.362	...	655 $\pm$ 40	8.70	5.9	40.00	2.11E37	779 $\pm$ 15
021004	2.33	...	470 $\pm$ 26	4.86	5.9	13.29	5.44E39	80 $\pm$ 38
030226	1.986	...	171 $\pm$ 23	8.46	9.7	3.43	1.24E38	97 $\pm$ 17
030329	0.169	80	10337 $\pm$ 33	4.86	2.2	5.68	1.58E38	68 $\pm$ 2.2
031203	0.105	0.3	828 $\pm$ 28	4.86	28.2	5.00	3.76E36	190 $\pm$ 30
050416A	0.65	0.47	485 $\pm$ 36	4.86	52.9	5.00	8.5E37	15 $\pm$ 2.7
050603	2.821	0.075	377 $\pm$ 53	8.46	29.4	5.00	9.89E38	344 $\pm$ 52
050820A	2.615	7.7	150 $\pm$ 31	8.46	3.7	5.07	3.62E38	246 $\pm$ 40
050904	6.29	0.09	76 $\pm$ 14	8.46	2.6	5.00	5.69E38	435 $\pm$ 90
051022	0.809	5	268 $\pm$ 32	8.46	4.8	5.00	8.57E37	510 $\pm$ 20
060218	0.033	0.36	245 $\pm$ 50	4.86	2.9	80.26	2.37E37	4.9 $\pm$ 0.3
070125	1.548	1.9	1028 $\pm$ 16	8.46	3.7	7.90	2.63E39	366 $\pm$ 51
071003	1.604	0.9	616 $\pm$ 57	8.46	33.0	5.00	6.7E38	799 $\pm$ 100
071010B	0.947	13	227 $\pm$ 114	4.86	2.5	5.00	8.12E37	52 $\pm$ 6
080603A	1.687	0.6	207 $\pm$ 26	8.46	6.4	5.00	2.45E38	85 $\pm$ 30
090313	3.375	2.5	435 $\pm$ 22	8.46	1.9	3.08	5.58E38	...
090323	3.57	2.0	243 $\pm$ 13	8.46	2.1	4.63	7.61E38	416 $\pm$ 76
090328	0.736	2.1	686 $\pm$ 26	8.46	3.4	8.85	5.74E38	592 $\pm$ 237
090423	8.26	0.1	50 $\pm$ 11	8.46	9.3	9.64	1.9E39	49 $\pm$ 3.8
090424	0.544	2.3	236 $\pm$ 37	8.46	3.6	6.33	5.62E37	162 $\pm$ 3.4
090715B	3	1.2	191 $\pm$ 36	8.46	3.4	1.39	4.23E37	134 $\pm$ 40
090902B	1.883	4.4	84 $\pm$ 16	8.46	2.3	5.00	1.19E38	775 $\pm$ 11
091020	1.71	1.1	399 $\pm$ 21	8.46	4.9	5.00	4.82E38	187 $\pm$ 34
100414A	1.368	2.0	524 $\pm$ 19	8.46	4.0	5.00	4.35E38	572 $\pm$ 30
100418A	0.62	1.8	522 $\pm$ 83	4.86	3.4	8.43	2.37E38	25 $\pm$ 30
100814A	1.44	0.45	496 $\pm$ 24	4.50	43.4	5.00	3.65E38	128 $\pm$ 23
991216	1.02	4.6	126	4.86	5.3	4.44	4.07E37	318 $\pm$ 30
000210	0.85	...	93	8.46	...	5.02	3.29E37	408 $\pm$ 14
050401	2.898	0.8	122	8.46	8.6	5.00	3.33E38	118 $\pm$ 18
050525A	0.606	0.29	164	8.46	4.6	5.00	3.01E37	81.2 $\pm$ 1.4
050730	3.968	1.5	212	8.46	1.5	5.00	8.92E38	104 $\pm$ 20
050824	0.83	0.8	152	8.46	8.4	5.00	5.1E37	15 $\pm$ 2
060418	1.49	1.5	216	8.46	1.8	5.00	2.07E38	230 $\pm$ 20
071020	2.146	1.0	141	8.46	4.0	5.00	2.45E38	322 $\pm$ 65
100901A	1.408	7.0	331	4.50	1.7	5.72	3.07E38	...
100906A	1.727	2.5	215	8.46	1.0	2.88	8.77E37	...




Fig. 1.— Correlation between radio luminosity at 5 GHz and peak synchrotron frequency for GRBs and blazars. The solid line shows the best linear fit. The 1σ dispersion is marked by the two dashed lines.




Fig. 2.— Correlation between radio luminosity at 5 GHz and spectral slope α_{RX} for GRBs and blazars. The solid line shows the best linear fit. The 1σ dispersion is marked by the two dashed lines.